

ORDINARY ALGEBRAIC CURVES WITH MANY AUTOMORPHISMS IN POSITIVE CHARACTERISTIC

GÁBOR KORCHMÁROS AND MARIA MONTANUCCI

Keywords: Algebraic curves, algebraic function fields, positive characteristic, automorphism groups.

Subject classifications: 14H37, 14H05.

ABSTRACT. Let $\mathcal{X}$ be an ordinary (projective, geometrically irreducible, nonsingular) algebraic curve of genus $g(\mathcal{X}) \geq 2$ defined over an algebraically closed field $\mathbb{K}$ of odd characteristic p . Let $\text{Aut}(\mathcal{X})$ be the group of all automorphisms of $\mathcal{X}$ which fix $\mathbb{K}$ element-wise. For any solvable subgroup G of $\text{Aut}(\mathcal{X})$ we prove that $|G| \leq 34(g(\mathcal{X}) + 1)^{3/2}$. There are known curves attaining this bound up to the constant 34. For p odd, our result improves the classical Nakajima bound $|G| \leq 84(g(\mathcal{X}) - 1)g(\mathcal{X})$, see [8], and, for solvable groups G , the Gunby-Smith-Yuan bound $|G| \leq 6(g(\mathcal{X})^2 + 12\sqrt{21}g(\mathcal{X})^{3/2})$ where $g(\mathcal{X}) > cp^2$ for some positive constant c ; see [2].

1. INTRODUCTION

In this paper, $\mathcal{X}$ stands for a (projective, geometrically irreducible, nonsingular) algebraic curve of genus $g(\mathcal{X}) \geq 2$ defined over an algebraically closed field $\mathbb{K}$ of odd characteristic p . Let $\text{Aut}(\mathcal{X})$ be the group of all automorphisms of $\mathcal{X}$ which fix $\mathbb{K}$ element-wise. The assumption $g(\mathcal{X}) \geq 2$ ensures that $\text{Aut}(\mathcal{X})$ is finite. However the classical Hurwitz bound $|\text{Aut}(\mathcal{X})| \leq 84(g(\mathcal{X}) - 1)$ for complex curves fails in positive characteristic, and there exist four families of curves satisfying $|\text{Aut}(\mathcal{X})| \geq 8g^3(\mathcal{X})$; see [11], Henn [4], and also [3, Section 11.12]. Each of them has p -rank $\gamma(\mathcal{X})$ (equivalently, its Hasse-Witt invariant) equal to zero; see for instance [1]. On the other hand, if $\mathcal{X}$ is ordinary, i.e. $g(\mathcal{X}) = \gamma(\mathcal{X})$, Guralnik and Zieve announced in 2004, as reported in [2, 5], that for odd p there exists a sharper bound, namely $|\text{Aut}(\mathcal{X})| \leq c_p g(\mathcal{X})^{8/5}$ with some constant c_p depending on p . It should be noticed that no proof of this sharper bound is available in the literature. In this paper, we concern with solvable automorphism groups G of an ordinary curve $\mathcal{X}$, and for odd p we prove the even sharper bound:

Theorem 1.1. *Let $\mathcal{X}$ be an algebraic curve of genus $g(\mathcal{X}) \geq 2$ defined over an algebraically closed field $\mathbb{K}$ of odd characteristic p . If $\mathcal{X}$ is ordinary and G is a solvable subgroup of $\text{Aut}(\mathcal{X})$ then*

$$(1) \quad |G| \leq 34(g(\mathcal{X}) + 1)^{3/2}.$$

For odd p , our result provides an improvement on the classical Nakajima bound $|G| \leq 84(g(\mathcal{X}) - 1)g(\mathcal{X})$, see [8], and, for solvable groups, on the recent Gunby-Smith-Yuan bound $|G| \leq 6(g(\mathcal{X})^2 + 12\sqrt{21}g(\mathcal{X})^{3/2})$ proven in [2] under the hypothesis that $g(\mathcal{X}) > cp^2$ for some positive constant c .

The following example is due to Stichtenoth and it shows that (1) is the best possible bound apart from the constant c . Let $\mathbb{F}_q$ be a finite field of order $q = p^h$ and let $\mathbb{K} = \overline{\mathbb{F}_q}$ denote its algebraic closure. For a positive integer m prime to p , let $\mathcal{Y}$ be the irreducible curve with affine equation

$$(2) \quad y^q + y = x^m + \frac{1}{x^m},$$

and $F = \mathbb{K}(\mathcal{Y})$ its function field. Let $t = x^{m(q-1)}$. The extension $F|\mathbb{K}(t)$ is a non-Galois extension as the Galois closure of F with respect to H is the function field $\mathbb{K}(x, y, z)$ where x, y, z are linked by (2) and $z^q + z = x^m$. Furthermore, $\mathfrak{g}(\mathcal{Y}) = (q-1)(qm-1)$, $\gamma(\mathcal{Y}) = (q-1)^2$ and $\text{Aut}(\mathcal{Y})$ contains a subgroup $Q \rtimes U$ of index 2 where Q is an elementary abelian normal subgroup of order q^2 and the complement U is a cyclic group of order $m(q-1)$. If $m = 1$ then $\mathcal{Y}$ is an ordinary curve, and in this case $2\mathfrak{g}(\mathcal{Y})^{3/2} = 2(q-1)^3 < 2q^2(q-1)$ which shows indeed that (1) is sharp up to the constant c .

2. BACKGROUND AND PRELIMINARY RESULTS

For a subgroup G of $\text{Aut}(\mathcal{X})$, let $\bar{\mathcal{X}}$ denote a non-singular model of $\mathbb{K}(\mathcal{X})^G$, that is, a (projective non-singular geometrically irreducible) algebraic curve with function field $\mathbb{K}(\mathcal{X})^G$, where $\mathbb{K}(\mathcal{X})^G$ consists of all elements of $\mathbb{K}(\mathcal{X})$ fixed by every element in G . Usually, $\bar{\mathcal{X}}$ is called the quotient curve of $\mathcal{X}$ by G and denoted by $\mathcal{X}/G$. The field extension $\mathbb{K}(\mathcal{X})|\mathbb{K}(\mathcal{X})^G$ is Galois of degree $|G|$.

Since our approach is mostly group theoretical, we prefer to use notation and terminology from Group theory rather than from Function field theory.

Let Φ be the cover of $\mathcal{X}|\bar{\mathcal{X}}$ where $\bar{\mathcal{X}} = \mathcal{X}/G$. A point $P \in \mathcal{X}$ is a ramification point of G if the stabilizer G_P of P in G is nontrivial; the ramification index e_P is $|G_P|$; a point $\bar{Q} \in \bar{\mathcal{X}}$ is a branch point of G if there is a ramification point $P \in \mathcal{X}$ such that $\Phi(P) = \bar{Q}$; the ramification (branch) locus of G is the set of all ramification (branch) points. The G -orbit of $P \in \mathcal{X}$ is the subset $o = \{R \mid R = g(P), g \in G\}$ of $\mathcal{X}$, and it is *long* if $|o| = |G|$, otherwise o is *short*. For a point $\bar{Q}$, the G -orbit o lying over $\bar{Q}$ consists of all points $P \in \mathcal{X}$ such that $\Phi(P) = \bar{Q}$. If $P \in o$ then $|o| = |G|/|G_P|$ and hence $\bar{Q}$ is a branch point if and only if o is a short G -orbit. It may be that G has no short orbits. This is the case if and only if every non-trivial element in G is fixed-point-free on $\mathcal{X}$, that is, the cover Φ is unramified. On the other hand, G has a finite number of short orbits. For a non-negative integer i , the i -th ramification group of $\mathcal{X}$ at P is denoted by $G_P^{(i)}$ (or $G_i(P)$ as in [10, Chapter IV]) and defined to be

$$G_P^{(i)} = \{g \mid \text{ord}_P(g(t) - t) \geq i + 1, g \in G_P\},$$

where t is a uniformizing element (local parameter) at P . Here $G_P^{(0)} = G_P$.

Let $\bar{\mathfrak{g}}$ be the genus of the quotient curve $\bar{\mathcal{X}} = \mathcal{X}/G$. The Hurwitz genus formula gives the following equation

$$(3) \quad 2\bar{\mathfrak{g}} - 2 = |G|(2\bar{\mathfrak{g}} - 2) + \sum_{P \in \mathcal{X}} d_P.$$

where the different d_P at P is given by

$$(4) \quad d_P = \sum_{i \geq 0} (|G_P^{(i)}| - 1).$$

Let γ be the p -rank of $\mathcal{X}$, and let $\bar{\gamma}$ be the p -rank of the quotient curve $\bar{\mathcal{X}} = \mathcal{X}/S$. The Deuring-Shafarevich formula, see [12] or [3, Theorem 11.62], states that

$$(5) \quad \gamma - 1 = |G|(\bar{\gamma} - 1) + \sum_{i=1}^k (|G| - \ell_i)$$

where $\ell_1, \dots, \ell_k$ are the sizes of the short orbits of G .

A subgroup of $\text{Aut}(\mathcal{X})$ is a prime to p group (or a p' -subgroup) if its order is prime to p . A subgroup G of $\text{Aut}(\mathcal{X})$ is *tame* if the 1-point stabilizer of any point in G is p' -group. Otherwise, G is *non-tame* (or *wild*). Obviously, every p' -subgroup of $\text{Aut}(\mathcal{X})$ is tame, but the converse is not always true. By a theorem of Stichtenoth, see [3, Theorem 11.56], if $|G| > 84(\mathfrak{g}(\mathcal{X}) - 1)$ then G is non-tame. An orbit o of G is *tame* if G_P is a p' -group for $P \in o$. The stabilizer G_P of a point $P \in \mathcal{X}$ in G is a semidirect product $G_P = Q_P \rtimes U$ where the normal subgroup Q_P is a p -group while the complement U is a cyclic prime to p group; see [3,

Theorem 11.49]. By a theorem of Serre, if $\mathcal{X}$ is an ordinary curve then Q_P is elementary abelian and no nontrivial element of U commutes with a nontrivial element of Q_P ; see [3, Lemma 11.75]. In particular, $|U|$ divides $|Q_P| - 1$, see [8, Proposition 1], and G_P is not abelian when either Q_P or U is nontrivial.

The following two lemmas of independent interest play a role in our proof of Theorem 1.1.

Lemma 2.1. *Let $\mathcal{X}$ be an ordinary algebraic curve of genus $\mathfrak{g}(\mathcal{X}) \geq 2$ defined over an algebraically closed field $\mathbb{K}$ of odd characteristic p . Let H be a solvable automorphism group of $\text{Aut}(\mathcal{X})$ containing a normal p -subgroup Q such that $|Q|$ and $[H : Q]$ are coprime. Suppose that a complement U of Q in H is abelian and that*

$$(6) \quad |H| > \begin{cases} 18(\mathfrak{g} - 1) & \text{for } |U| = 3, \\ 12(\mathfrak{g} - 1) & \text{otherwise.} \end{cases}$$

Then U is cyclic, and the quotient curve $\bar{\mathcal{X}} = \mathcal{X}/Q$ is rational. Furthermore, Q has exactly two (non-tame) short orbits, say Ω_1, Ω_2 . They are also the only short orbits of H , and $\mathfrak{g}(\mathcal{X}) = |Q| - (|\Omega_1| + |\Omega_2|) + 1$.

Proof. From the Schur-Zassenhaus theorem [7, Corollary 7.5], $H = Q \rtimes U$. Set $|Q| = p^k$, $|U| = u$. Then $p \nmid u$. Furthermore, if $u = 2$ then $|H| = 2|Q| > 9\mathfrak{g}(\mathcal{X})$ whence $|Q| > 4.5\mathfrak{g}(\mathcal{X})$. From Nakajima's bound $\mathcal{X}$, see [3, Theorem 11.84], has zero p -rank. Therefore $u \geq 3$.

Three cases are treated separately according as the quotient curve $\bar{\mathcal{X}} = \mathcal{X}/Q$ has genus $\bar{\mathfrak{g}}$ at least 2, or $\bar{\mathcal{X}}$ is elliptic, or rational.

If $\mathfrak{g}(\bar{\mathcal{X}}) \geq 2$, then $\text{Aut}(\bar{\mathcal{X}})$ has a subgroup isomorphic to U , and [3, Theorem 11.79] yields $4\mathfrak{g}(\bar{\mathcal{X}}) + 4 \geq |U|$. Furthermore, from the Hurwitz genus formula applied to Q , $\mathfrak{g} - 1 \geq |Q|(\mathfrak{g}(\bar{\mathcal{X}}) - 1)$. Therefore, if $c = 12$ or $c = 18$, according as $|U| > 3$ or $|U| = 3$,

$$(4\mathfrak{g}(\bar{\mathcal{X}}) + 4)|Q| \geq |U||Q| = |H| > c(\mathfrak{g} - 1) = 12|Q|(\mathfrak{g}(\bar{\mathcal{X}}) - 1),$$

whence

$$c < 4 \frac{\mathfrak{g}(\bar{\mathcal{X}}) + 1}{\mathfrak{g}(\bar{\mathcal{X}}) - 1} \leq 12,$$

a contradiction.

If $\bar{\mathcal{X}}$ is elliptic, then the cover $\mathcal{X}|\bar{\mathcal{X}}$ ramifies, otherwise $\mathcal{X}$ itself would be elliptic. Thus, Q has some short orbits. Take one of them together with its images $o_1, \dots, o_{u_1}$ under the action of H . Since Q is a normal subgroup of H , $o = o_1 \cup \dots \cup o_{u_1}$ is a H -orbit of size $u_1 p^v$ where $p^v = |o_1| = \dots = |o_{u_1}|$. Equivalently, the stabilizer of a point $P \in o$ has order $p^{k-v} u / u_1$, and it is the semidirect product $Q_1 \rtimes U_1$ where $|Q_1| = p^{k-v}$ and $|U_1| = u / u_1$ for a subgroup Q_1 of Q and U_1 of U respectively. The point $\bar{P}$ lying under P in the cover $\mathcal{X}|\bar{\mathcal{X}}$ is fixed by the factor group $\bar{U}_1 = U_1 Q / Q$. Since $\bar{\mathcal{X}}$ is elliptic, [3, Theorem 11.94] implies $|\bar{U}_1| \leq 12$ for $p = 3$ and $|\bar{U}_1| \leq 6$ for $p > 3$. As $\bar{U}_1 \cong U_1$, this yields the same bound for $|U_1|$, that is, $u \leq 4u_1$ for $p = 3$ and $u \leq 6u_1$ for $p > 3$. Furthermore, $d_P \geq 2(p^{k-v} - 1) \geq \frac{4}{3}p^{k-v}$. From the Hurwitz genus formula applied to Q , if $p = 3$ then

$$2\mathfrak{g} - 2 \geq 3^v u_1 d_P \geq 3^v u_1 (\frac{4}{3} 3^{k-v}) \geq \frac{4}{3} 3^k u_1 \geq \frac{1}{3} 3^k u = \frac{1}{3} |Q| |U| = \frac{1}{3} |H|,$$

while for $p > 3$,

$$2\mathfrak{g} - 2 \geq p^v u_1 d_P \geq p^v u_1 (\frac{4}{3} p^{k-v}) \geq \frac{4}{3} p^k u_1 \geq \frac{2}{9} p^k u = \frac{2}{9} |Q| |U| = \frac{2}{9} |H|,$$

But this contradicts (6).

If $\bar{\mathcal{X}}$ is rational, then Q has at least one short orbit. Furthermore, $\bar{U} = UQ/Q$ is isomorphic to a subgroup of $PGL(2, \mathbb{K}) \cong \text{Aut}(\bar{\mathcal{X}})$. Since $U \cong \bar{U}$, the classification of finite subgroups of $PGL(2, \mathbb{K})$, see [13], shows that U is cyclic, $\bar{U}$ fixes two points $\bar{P}_0$ and $\bar{P}_\infty$ but no nontrivial element in $\bar{U}$ fixes a point other than $\bar{P}_0$ and $\bar{P}_\infty$. Let o_∞ and o_0 be the Q -orbits lying over $\bar{P}_0$ and $\bar{P}_\infty$, respectively. Obviously, o_∞ and o_0 are short orbits of H . We show that Q has at most two short orbits, the candidates being o_∞ and o_0 . By absurd, there is a Q -orbit o of size p^m with $m < k$ which lies over a point $\bar{P} \in \bar{\mathcal{X}}$ different from both $\bar{P}_0$ and $\bar{P}_\infty$.

Since the orbit of $\bar{P}$ in $\bar{U}$ has length u , then the H -orbit of a point $P \in o$ has length up^m . If $u > 3$, the Hurwitz genus formula applied to Q gives

$$2\mathfrak{g} - 2 \geq -2p^k + up^m(p^{k-m} - 1) \geq -2p^k + up^m \frac{2}{3}p^{k-m} \geq -2p^k + \frac{2}{3}up^k \geq \frac{2}{3}(u-3)p^k \geq \frac{1}{6}up^k = \frac{1}{6}|H|,$$

a contradiction with $|H| > 12(\mathfrak{g} - 1)$. If $u = 3$ then $p > 3$, and hence

$$2\mathfrak{g} - 2 \geq -2p^k + 3p^m(p^{k-m} - 1) = p^k - 3p^m > \frac{1}{3}p^k,$$

whence $|H| = 3p^k < 18(\mathfrak{g} - 1)$, a contradiction with (6). This proves that H has exactly two short orbits. Assume that Q has two short orbits. They are o_∞ and o_0 . If their lengths are p^a and p^b with $a, b < k$, the Deuring-Shafarevich formula applied to Q gives

$$\mathfrak{g}(\mathcal{X}) - 1 = \gamma(\mathcal{X}) - 1 = -p^k + (p^k - p^a) + p^k - p^b$$

whence $\mathfrak{g}(\mathcal{X}) = p^k - (p^a + p^b) + 1 > 0$. The same argument shows that if Q has just one short orbit, then $\gamma(\mathcal{X}) = 0$, a contradiction. $\square$

Lemma 2.2. *Let N be an automorphism group of an algebraic curve of even genus such that $|N|$ is even. Then any 2-subgroup of N has a cyclic subgroup of index 2.*

Proof. Let U be a subgroup of $\text{Aut}(\mathcal{X})$ of order $d = 2^u \geq 2$, and $\bar{\mathcal{X}} = \mathcal{X}/N$ the arising quotient curve. From the Hurwitz genus formula applied to U ,

$$2\mathfrak{g}(\mathcal{X}) - 2 = 2(p-2)p^{n-1} = 2^u(2\mathfrak{g}\bar{\mathcal{X}} - 2) + \sum_{i=1}^m (2^u - \ell_i)$$

where $\ell_1, \dots, \ell_m$ are the short orbits of U on $\mathcal{X}$. Since $2(p-2)p^{n-1} \equiv 2 \pmod{4}$ while $2^u(2\mathfrak{g}\bar{\mathcal{X}} - 2) \equiv 0 \pmod{4}$, some ℓ_i ($1 \leq i \leq m$) must be either 1 or 2. Therefore, U or a subgroup of U of index 2 fixes a point of $\mathcal{X}$ and hence is cyclic. $\square$

3. THE PROOF OF THEOREM 1.1

The assertion holds for $\mathfrak{g}(\mathcal{X}) = 2$, as $|G| \leq 48$ for any solvable automorphism group G of a genus two curve; see [3, Proposition 11.99]. For $\mathfrak{g}(\mathcal{X}) > 2$, $\mathcal{X}$ is taken by absurd for a minimal counterexample with respect the genera so that for any solvable subgroup of $\text{Aut}(\mathcal{X})$ of an ordinary curve $\bar{\mathcal{X}}$ of genus $\mathfrak{g}(\bar{\mathcal{X}}) \geq 2$ we have $|\bar{G}| \leq 34(\mathfrak{g} + 1)^{3/2}$. Two cases are treated separately.

3.1. Case I: G contains a minimal normal p -subgroup.

Proposition 3.1. *Let $\mathcal{X}$ be an ordinary algebraic curve of genus $\mathfrak{g}$ defined over an algebraically closed field $\mathbb{K}$ of odd characteristic $p > 0$. If G is a solvable subgroup of $\text{Aut}(\mathcal{X})$ containing a minimal normal p -subgroup N , then $|G| \leq 34(\mathfrak{g} + 1)^{3/2}$.*

Proof. Take the largest normal p -subgroup Q of G . Let $\bar{\mathcal{X}}$ be the quotient curve of $\mathcal{X}$ with respect to Q and let $\bar{G} = G/Q$. The quotient group $\bar{G}$ is a subgroup of $\text{Aut}(\bar{\mathcal{X}})$ and it has no normal p -subgroup, otherwise G would have a normal p -subgroup properly containing Q . For $\bar{\mathfrak{g}} = \mathfrak{g}(\bar{\mathcal{X}})$ three cases may occur, namely $\bar{\mathfrak{g}} \geq 2$, $\bar{\mathfrak{g}} = 1$ or $\bar{\mathfrak{g}} = 0$. If $\bar{\mathfrak{g}} \geq 2$, from the Hurwitz genus formula,

$$2\mathfrak{g} - 2 \geq |Q|(2\bar{\mathfrak{g}} - 2) = \frac{|G|}{|\bar{G}|}(2\bar{\mathfrak{g}} - 2)$$

whence $|\bar{G}| > 34(\bar{\mathfrak{g}} + 1)^{3/2}$. Since $\bar{\mathcal{X}}$ is still ordinary, this contradicts our choice of $\mathcal{X}$ to be a minimal counterexample. If $\bar{\mathfrak{g}} = 1$ then the cover $\mathbb{K}(\mathcal{X})|\mathbb{K}(\bar{\mathcal{X}})$ ramifies. Take short orbit Δ of Q . Let Γ be the non-tame short orbit of G that contains Δ . Since Q is normal in G , the orbit Γ partitions into short orbits

of Q whose components have the same length which is equal to $|\Delta|$. Let k be the number of the Q -orbits contained in Γ . Then,

$$|G_P| = \frac{|G|}{k|\Delta|},$$

holds for every $P \in \Gamma$. Moreover, the quotient group $G_P Q/Q$ fixes a place on $\bar{\mathcal{X}}$. Now, from [3, Theorem 11.94 (ii)],

$$\frac{|G_P Q|}{|Q|} = \frac{|G_P|}{|G_P \cap Q|} = \frac{|G_P|}{|Q_P|} \leq 12.$$

From this together with the Hurwitz genus formula and [8, Theorem 2 (i)],

$$2g - 2 \geq 2k|\Delta|(|Q_P| - 1) \geq 2k|\Delta| \frac{|Q_P|}{2} \geq \frac{k|\Delta||G_P|}{12} = \frac{|G|}{12},$$

which contradicts our hypothesis $|G| > 34(g+1)^{3/2}$.

It turns out that $\bar{\mathcal{X}}$ is rational. Therefore $\bar{G}$ is isomorphic to a subgroup of $PGL(2, \mathbb{K})$ which contains no normal p -subgroup. From the classification of finite subgroups of $PGL(2, \mathbb{K})$, see [13], $\bar{G}$ is a prime to p subgroup which is either cyclic, or dihedral, or isomorphic to one of the groups $\text{Alt}_4, \text{Sym}_4$. In all cases, $\bar{G}$ has a cyclic subgroup U of index ≤ 6 and of order distinct from 3. We may dismiss all cases but the cyclic one assuming that $G = Q \rtimes U$ with $|G| \geq \frac{5}{3}(g(\mathcal{X}) + 1)^{3/2}$. Then $|G| > 12(g-1)$. Therefore, Lemma 2.1 applies to G . Thus, Q has exactly two (non-tame) orbits, say Ω_1 and Ω_2 , and they are also the only short orbits of G . More precisely,

$$(7) \quad \gamma - 1 = |Q| - (|\Omega_1| + |\Omega_2|).$$

We may also observe that G_P with $P \in \Omega_1$ contains a subgroup V isomorphic to U . In fact, $|Q||U| = |G| = |G_P||\Omega_1| = |Q_P \rtimes V||\Omega_1| = |V||Q_P||\Omega_1|$ with a prime to p subgroup V fixing P , whence $|U| = |V|$. Since V is cyclic the claims follows.

We go on with the case where both Ω_1 and Ω_2 are nontrivial, that is, their lengths are at least 2.

Assume that Q is non-abelian and look at the action of its center $Z(Q)$ on $\mathcal{X}$. Since $Z(Q)$ is a nontrivial normal subgroup of G , arguing as before we get that the factor group $\mathcal{X}/Z(Q)$ is rational, and hence the Galois cover $\mathcal{X}/(\mathcal{X}/Z(Q))$ ramifies at some points. In other words, there is a point $P \in \Omega_1$ (or $R \in \Omega_2$) such that some nontrivial subgroup T of $Z(Q)$ fixes P (or Q). Suppose that the former case occurs. Since Ω_1 is a Q -orbit, T fixes Ω_1 pointwise.

The group G induces a permutation group on Ω_1 and let M_1 be the kernel of this permutation representation. Obviously, T is a nontrivial p -subgroup of M_1 . Therefore M contains some but not all elements from Q . Since both M_1 and Q are normal subgroups of G , $N = M_1 \cap Q$ is a nontrivial normal p -subgroup of G . From (ii), the quotient curve $\tilde{\mathcal{X}} = \mathcal{X}/N$ is rational, and hence the factor group $\tilde{G} = G/N$ is isomorphic to a subgroup of $PGL(2, \mathbb{K})$. Since $1 \not\leq N \leq Q$, the order of $\tilde{G}$ is divisible by p . From the classification of subgroups of $PGL(2, \mathbb{K})$, see [13], $\tilde{G} = \tilde{Q} \rtimes \tilde{U}$ where $\tilde{Q}$ is an elementary abelian p -group of order q and $\tilde{U} \cong UN/N \cong U$ with $|\tilde{U}| = |U|$ is a divisor of $q-1$.

This shows that Q acts on Ω_1 as an abelian transitive permutation group. Obviously this holds true when Q is abelian. Therefore, the action of Q on Ω_1 is sharply transitive. In terms of 1-point stabilizers of Q on Ω_1 , we have $Q_P = Q_{P'}$ for any $P, P' \in \Omega_1$. Moreover, $Q_P = N$, and hence Q_P is a normal subgroup of G .

Furthermore, since $\mathcal{X}$ is an ordinary curve, Q_P is an elementary abelian group by [8, Theorem 2 (i)] and [3, Theorem 11.74 (iii)].

The quotient curve $\mathcal{X}/Q_P$ is rational and its automorphism group contains the factor group Q/Q_P . Hence, exactly one of the Q_P -orbits is preserved by Q . Since Ω_1 is a Q -orbit consisting of fixed points of Q_P , Ω_2 must be a Q_P -orbit. Similarly, if $Z(Q) \neq Q_P$, the factor group $Z(Q)Q_P/Q_P$ is an automorphism group of $\mathcal{X}/Q_P$ and hence exactly one of the Q_P -orbits is preserved by $Z(Q)$. Either $Z(Q)$ fixes a point in Ω_1 but then $Z(Q) = Q_P$, or Ω_2 is a $Z(Q)$ -orbit. This shows that either $Z(Q) = Q_P$ or $Z(G)$ acts transitively on Ω_2 .

Two cases arise according as Q_P is sharply transitive and faithful on Ω_2 or some nontrivial element in Q_P fixes Ω_2 pointwise.

If some nontrivial element in Q_P fixes Ω_2 pointwise then the kernel M_2 of the permutation representation of H on Ω_2 contains a nontrivial p -subgroup. Hence the above results extends from Ω_1 to Ω_2 , and Q_R is a normal subgroup of Q .

If Q_P is (sharply) transitive on Ω_2 then the abelian group $Z(Q)Q_P$ acts on Ω_2 as a sharply transitive permutation group, as well. Hence either $Z(Q) = Q_P$, or as before M_2 contains a nontrivial p -subgroup, and Q_R is a normal subgroup of Q . In the former case, $Q = Q_P Q_R$ and $Q_R \cap Q_P = \{1\}$, and $Z(Q) = Q_P$ yields that

$$(8) \quad Q = Q_P \times Q_R.$$

This shows that Q is abelian, and hence $|Q| \leq 4g + 4$. Also, either $|Q_P|$ (or $|Q_R|$) is at most $\sqrt{4g + 4}$. Since $\mathcal{X}$ is an ordinary curve, the second ramification group $G_P^{(2)}$ at $P \in \Omega_1$ is trivial. For $G_P = Q_P \rtimes V$, [3, Lemma 11.81] gives $|U| = |V| \leq |Q_P| - 1$. Hence $|U| < |Q_P| \leq \sqrt{|Q|} \leq \sqrt{4g + 4}$ whence

$$(9) \quad |G| = |U||Q| \leq 8(g + 1)^{3/2}.$$

If Q_R is a normal subgroup, take a point R from Ω_2 , and look at the subgroup $Q_{P,R}$ of Q_P fixing R . Actually, we prove that that either $Q_{P,R} = Q_P$ or $Q_{P,R}$ is trivial. Suppose that $Q_{P,R} \neq \{1\}$. Since $Q_{P,R} = Q_P \cap Q_R$ and both Q_P and Q_R are normal subgroups of G , the same holds for $Q_{P,R}$. By (ii), the quotient curve $\mathcal{X}/Q_{P,R}$ is rational and hence its automorphism group $Q/Q_{P,R}$ fixes exactly one point. Furthermore, each point in Ω_2 is totally ramified. Therefore, $Q_R = Q_{P,R}$, otherwise $Q_R/Q_{P,R}$ would fix any point lying under a point in Ω_1 in the cover $\mathcal{X}/(Q/Q_{P,R})$.

It turns out that either $Q_P = Q_R$ or $Q_P \cap Q_R = \{1\}$, whenever $P \in \Omega_1$ and $R \in \Omega_2$.

In the former case, from the Deuring-Shafarevič formula applied to Q_P ,

$$\gamma - 1 = -|Q_P| + |\Omega_1|(|Q_P| - 1) + |\Omega_2|(|Q_P| - 1) = -|Q_P| + |Q| - |\Omega_1| + |Q| - |\Omega_2|.$$

This together with (7) give $Q = Q_P$, a contradiction.

Therefore, the latter case must hold. Thus $Q = Q_P \times Q_R$ and Q_P (and also Q_R) is an elementary abelian group since is isomorphic to a p -subgroup of $PGL(2, \mathbb{K})$. Also, $|Q_P| = |Q_R| = \sqrt{|Q|}$. Since Q is abelian, this yields $|Q_P| \leq \sqrt{4g + 4}$. Now, the argument used after (8) can be employed to prove (9). This ends the proof in the case where both Ω_1 and Ω_2 are nontrivial.

Suppose next $\Omega_1 = \{P\}$ and $|\Omega_2| \geq 2$. Then G fixes P , and hence $G = Q \rtimes U$ with an elementary abelian p -group Q . Furthermore, G has a permutation representation on Ω_2 with kernel K . As Ω_2 is a short orbit of Q , the stabilizer Q_R of $R \in \Omega_2$ in Q is nontrivial. Since Q is abelian, this yields that K is nontrivial, and hence it is a nontrivial elementary abelian normal subgroup of G . In other words, Q is an r -dimensional vector space $V(r, p)$ over a finite field $\mathbb{F}_p$ with $|Q| = p^r$, the action of each nontrivial element of U by conjugacy is a nontrivial automorphism of $V(r, p)$, and K is a K -invariant subspace. By Maschke's theorem, see for instance [7, Theorem 6.1], K has a complementary U -invariant subspace. Therefore, Q has a subgroup M such that $Q = K \times M$, and M is a normal subgroup of G . Since $K \cap M = \{1\}$, and Ω_2 is an orbit of Q , this yields $|M| = |\Omega_2|$. The factor group G/M is an automorphism group of the quotient curve $\mathcal{X}/M$, and Q/M is a nontrivial p -subgroup of G/M whereas G/M fixes two points on $\mathcal{X}/M$. Therefore the quotient curve $\mathcal{X}/M$ is not rational since the 2-point stabilizer in the representation of $PGL(2, \mathbb{K})$ as an automorphism group of the rational function field is a prime to p (cyclic) group. We show that $\mathcal{X}/M$ is neither elliptic. From the Deuring-Shafarevič formula, $g(\mathcal{X}) - 1 = \gamma(\mathcal{X}) - 1 = -|Q| + 1 + |\Omega_2|$, and so $g(\mathcal{X})$ is even. Since M is a normal subgroup of odd order, $g(\mathcal{X}) \equiv 0 \pmod{2}$ yields that $g(\mathcal{X}/M) \equiv 0 \pmod{2}$. In particular, $g(\mathcal{X}/M) \neq 1$. Therefore, $g(\mathcal{X}/M) \geq 2$. At this point we may repeat our previous argument and prove $|G/M| > 34(g(\mathcal{X}/M) + 1)^{3/2}$. Again, a contradiction to our choice of $\mathcal{X}$ to be a minimal counterexample, which ends the proof in the case where just one of Ω_1 and Ω_2 is trivial.

We are left with the case where both short orbits of Q are trivial. Our goal is to prove a much stronger bound for this case, namely $|U| \leq 2$ whence

$$(10) \quad |G| \leq 2(g(\mathcal{X}) + 1).$$

We also show that if equality holds then $\mathcal{X}$ is a hyperelliptic curve with equation

$$(11) \quad f(U) = aT + b + cT^{-1}, \quad a, b, c \in \mathbb{K}^*,$$

where $f(U) \in \mathbb{K}[U]$ is an additive polynomial of degree $|Q|$.

Let $\Omega_1 = \{P_1\}$ and $\Omega_2 = \{P_2\}$. Then Q has two fixed points P_1 and P_2 but no nontrivial element in Q fixes a point of $\mathcal{X}$ other than P_1 and P_2 . From the Deuring-Shafarevič formula

$$(12) \quad g(\mathcal{X}) + 1 = \gamma(\mathcal{X}) + 1 = |Q|.$$

Therefore, $|U| \leq g(\mathcal{X})$. Actually, for our purpose, we need a stronger estimate, namely $|U| \leq 2$. To prove the latter bound, we use some ideas from Nakajima's paper [8]. regarding the Riemann-Roch spaces $\mathcal{L}(\mathbf{D})$ of certain divisors $\mathbf{D}$ of $\mathbb{K}(\mathcal{X})$. Our first step is to show

- (i) $\dim_{\mathbb{K}} \mathcal{L}((|Q| - 1)P_1) = 1$,
- (ii) $\dim_{\mathbb{K}} \mathcal{L}((|Q| - 1)P_1 + P_2) \geq 2$.

Let $\ell \geq 1$ be the smallest integer such that $\dim_{\mathbb{K}} \mathcal{L}(\ell P_1) = 2$, and take $x \in \mathcal{L}(\ell P_1)$ with $v_{P_1}(x) = -\ell$. As $Q = Q_{P_1}$, the Riemann-Roch space $\mathcal{L}(\ell P_1)$ contains all $c_\sigma = \sigma(x) - x$ with $\sigma \in Q$. This yields $c_\sigma \in \mathbb{K}$ by $v_{P_1}(c_\sigma) \geq -\ell + 1$ and our choice of ℓ to be minimal. Also, $Q = Q_{P_2}$ together with $v_{P_2}(x) \geq 0$ show $v_{P_2}(c_\sigma) \geq 1$. Therefore $c_\sigma = 0$ for all $\sigma \in Q$, that is, x is fixed by Q . From $\ell = [\mathbb{K}(\mathcal{X}) : \mathbb{K}(x)] = [\mathbb{K} : \mathbb{K}(\mathcal{X})^Q][\mathbb{K}(\mathcal{X})^Q : \mathbb{K}(x)]$ and $|Q| = [\mathbb{K} : \mathbb{K}(\mathcal{X})^Q]$, it turns out that ℓ is a multiple of $|Q|$. Thus $\ell > |Q| - 1$ whence (i) follows. From the Riemann-Roch theorem, $\dim_{\mathbb{K}} \mathcal{L}((|Q| - 1)P_1 + P_2) \geq |Q| - g + 1 = 2$ which proves (ii).

Let $d \geq 1$ be the smallest integer such that $\dim_{\mathbb{K}} \mathcal{L}(dP_1 + P_2) = 2$. From (ii)

$$(13) \quad d \leq |Q| - 1.$$

Let α be a generator of the cyclic group U . Since α fixes both points P_1 and P_2 , it acts on $\mathcal{L}(dP_1 + P_2)$ as a $\mathbb{K}$ -vector space automorphism $\bar{\alpha}$. If $\bar{\alpha}$ is trivial then $\alpha(u) = u$ for all $u \in \mathcal{L}(dP_1 + P_2)$. Suppose that $\bar{\alpha}$ is nontrivial. Since U is a prime to p cyclic group, $\bar{\alpha}$ has two distinct eigenspaces, so that $\mathcal{L}(dP_1 + P_2) = \mathbb{K} \oplus \mathbb{K}u$ where $u \in \mathcal{L}(dP_1 + P_2)$ is an eigenvector of $\bar{\alpha}$ with eigenvalue $\xi \in \mathbb{K}^*$ so that $\bar{\alpha}(u) = \xi u$ with $\xi^{|U|} = 1$. Therefore there is $u \in \mathcal{L}(dP_1 + P_2)$ with $u \neq 0$ such that $\alpha(u) = \xi u$ with $\xi^{|U|} = 1$. The pole divisor of u is

$$(14) \quad \operatorname{div}(u)_\infty = dP_1 + P_2.$$

Since $Q = Q_{P_1} = Q_{P_2}$, the Riemann-Roch space $\mathcal{L}(dP_1 + P_2)$ contains $\sigma(u)$ and hence it contains all

$$\theta_\sigma = \sigma(u) - u, \quad \sigma \in Q.$$

By our choice of d to be minimal, this yields $\theta_\sigma \in \mathbb{K}$, and then defines the map θ from Q into $\mathbb{K}$ that takes σ to θ_σ . More precisely, θ is a homomorphism from Q into the additive group $(\mathbb{K}, +)$ of $\mathbb{K}$ as the following computation shows:

$$\theta_{\sigma_1 \circ \sigma_2} = (\sigma_1 \circ \sigma_2)(u) - u = \sigma_1(\sigma_2(u) - u + u) - u = \sigma_1(\theta_{\sigma_2}) + \sigma_1(u) - u = \theta_{\sigma_2} + \theta_{\sigma_1} = \theta_{\sigma_1} + \theta_{\sigma_2}.$$

Also, θ is injective. In fact, if $\theta_{\sigma_0} = 0$ for some $\sigma_0 \in Q \setminus \{1\}$, then u is in the fixed field of σ_0 , which is impossible since $v_{P_2}(u) = -1$ whereas P_2 is totally ramified in the cover $\mathcal{X} | (\mathcal{X} / \langle \sigma_p \rangle)$. The image $\theta(Q)$ of θ is an additive subgroup of $\mathbb{K}$ of order $|Q|$. The smallest subfield of $\mathbb{K}$ containing $\theta(Q)$ is a finite field $\mathbb{F}_{p^m}$ and hence $\theta(Q)$ can be viewed as a linear subspace of $\mathbb{F}_{p^m}$ considered as a vector space over $\mathbb{F}_p$. Therefore the polynomial

$$(15) \quad f(U) = \prod_{\sigma \in Q} (U - \theta_\sigma)$$

is a linearized polynomial over $\mathbb{F}_p$; see [6, Section 4, Theorem 3.52]. In particular, $f(U)$ is an additive polynomial of degree $|Q|$; see also [9, Chapter V, § 5]. Also, $f(U)$ is separable as θ is injective. From (15), the pole divisor of $f(u) \in \mathbb{K}(\mathcal{X})$ is

$$(16) \quad \operatorname{div}(f(u))_\infty = |Q|(dP_1 + P_2).$$

For every $\sigma_0 \in Q$,

$$\sigma_0(f(u)) = \prod_{\sigma \in Q} (\sigma_0(u) - \theta_\sigma) = \prod_{\sigma \in Q} (u + \theta_{\sigma_0} - \theta_\sigma) = \prod_{\sigma \in Q} (u - \theta_{\sigma\sigma_0^{-1}}) = \prod_{\sigma \in Q} (u - \theta_\sigma) = f(u).$$

Thus $f(u) \in \mathbb{K}(\mathcal{X})^Q$. Furthermore, from $\alpha \in N_G(Q)$, for every $\sigma \in Q$ there is $\sigma' \in Q$ such that $\alpha\sigma = \sigma'\alpha$. Therefore

$$\alpha(f(u)) = \prod_{\sigma \in Q} (\alpha(\sigma(u) - u)) = \prod_{\sigma \in Q} (\alpha(\sigma(u)) - \xi u) = \prod_{\sigma \in Q} (\sigma'(\alpha(u)) - \xi u) = \prod_{\sigma \in Q} (\sigma'(\xi u) - \xi u) = \xi f(u).$$

This shows that if $R \in \mathcal{X}$ is a zero of $f(u)$ then $\operatorname{Supp}(\operatorname{div}(f(u)_0))$ contains the U -orbit of R of length $|U|$. Actually, since $\sigma(f(u)) = f(u)$ for $\sigma \in Q$, $\operatorname{Supp}(\operatorname{div}(f(u)_0))$ contains the G -orbit of R of length $|G| = |Q||U|$. This together with (16) give

$$(17) \quad |U|(d+1).$$

On the other hand, $\mathbb{K}(\mathcal{X})^Q$ is rational. Let $\bar{P}_1$ and $\bar{P}_2$ the points lying under P_1 and P_2 , respectively, and let $\bar{R}_1, \bar{R}_2, \dots, \bar{R}_k$ with $k = (d+1)/|U|$ be the points lying under the zeros of $f(u)$ in the cover $\mathcal{X}|(\mathcal{X}/Q)$. We may represent $\mathbb{K}(\mathcal{X})^Q$ as the projective line $\mathbb{K} \cup \{\infty\}$ over $\mathbb{K}$ so that $\bar{P}_1 = \infty$, $\bar{P}_1 = 0$ and $\bar{R}_i = t_i$ for $1 \leq i \leq k$. Let $g(t) = t^d + t^{-1} + h(t)$ where $h(t) \in \mathbb{K}[t]$ is a polynomial of degree $k = (d+1)/|U|$ whose roots are $r_1, \dots, r_k$. It turns out that $f(u), g(t) \in \mathbb{K}(\mathcal{X})$ have the same pole and zero divisors, and hence

$$(18) \quad cf(u) = t^d + t^{-1} + h(t), \quad c \in \mathbb{K}^*.$$

We prove that $\mathbb{K}(\mathcal{X}) = \mathbb{K}(u, t)$. From [12], see also [3, Remark 12.12], the polynomial $cTf(X) - T^{d+1} - 1 - h(T)T$ is irreducible, and the plane curve $\mathcal{C}$ has genus $\mathfrak{g}(\mathcal{C}) = \frac{1}{2}(q-1)(d+1)$. Comparison with (12) shows $\mathbb{K}(\mathcal{X}) = \mathcal{C}$ and $d = 1$ whence $|U| \leq 2$. If equality holds then $\deg h(T) = 1$ and $\mathcal{X}$ is a hyperelliptic curve with equation (11). $\square$

3.2. Case II: G contains no minimal normal p -subgroup.

Proposition 3.2. *Let $\mathcal{X}$ be an ordinary algebraic curve of genus $\mathfrak{g}$ defined over a field $\mathbb{K}$ of odd characteristic $p > 0$. If G is a solvable subgroup of $\operatorname{Aut}(\mathcal{X})$ with a minimal normal subgroup N satisfying Case II, then $|G| \leq 34(\mathfrak{g}(\mathcal{X}) + 1)^{3/2}$.*

Proof. From Proposition 3.1, G contains no nontrivial normal p -subgroup. The factor group $\bar{G} = G/N$ is a subgroup of $\operatorname{Aut}(\mathcal{X})$ where $\bar{\mathcal{X}} = \mathcal{X}/N$. Furthermore, $|N| \leq 4\mathfrak{g}(\mathcal{X}) + 4$ as N is abelian; see [3, Theorem 11.79]. Arguing as in Section 3.1, we see that $\bar{\mathcal{X}}$ is either rational or elliptic, that is, $\mathfrak{g}(\bar{\mathcal{X}}) \geq 2$ is impossible.

A detailed description of the arguments is given below. Assume that $\mathcal{X}$ is a minimal counterexample with respect to the genus. Since $\mathcal{X}$ is ordinary, any p -subgroup S of G is an elementary abelian group and it has a trivial second ramification group at any point $\mathcal{X}$. The latter property remains true when $\mathcal{X}$ is replaced by $\bar{\mathcal{X}}$. To show this claim, take $\bar{P} \in \bar{\mathcal{X}}$ and let $\bar{S}_{\bar{P}}$ the subgroup of the factor group $\bar{S} = SN/N$ fixing $\bar{P}$. Since $p \nmid |N|$ there is a point $P \in \mathcal{X}$ lying over $\bar{P}$ which is fixed by S . Hence the stabilizer S_P of P in S is a nontrivial normal subgroup of G_P . Since N is a normal subgroup in G , so is N_P in G_P . This yields that the product $N_P S_P$ is actually a direct product. Therefore N_P is trivial, otherwise the second ramification group of S in P is nontrivial by a result of Serre; see [3, Theorem 11.75]. Actually, N may be assumed to be the largest normal subgroup N_1 of G whose order is prime to p . If the quotient curve $\mathcal{X}_1 = \mathcal{X}/N_1$ is neither rational, nor elliptic then its $\mathbb{K}$ -automorphism group $G_1 = G/N_1$ has order bigger than $34(\mathfrak{g}(\mathcal{X}_1) + 1)^{3/2}$, by the Hurwitz genus formula applied to N_1 . Since G and hence G_1 is solvable, G_1 has a minimal normal d -subgroup where d must

be equal to p by the choice of N_1 to be the largest normal, prime to p subgroup of G . Take the largest normal p -subgroup N_2 of G_1 . Observe that $N_2 \not\leq G_1$, otherwise G_1 is an (elementary) abelian group of order bigger than $34(\mathbf{g}(\mathcal{X}_1) + 1)^{3/2}$ contradicting the bound $4(\mathbf{g}(\mathcal{X}_1) + 4)$; see [3, Theorem 11.79]. Now, define $\mathcal{X}_2$ to be the quotient curve $\mathcal{X}_1/N_2$. Since the second ramification group of N_1 at any point of $\mathcal{X}_1$ is trivial, the Hurwitz genus formula together with the Deuring-Shafarevich formula give $\mathbf{g}(\mathcal{X}_1) - \gamma(\mathcal{X}_1) = |N_2|(\mathbf{g}(\mathcal{X}_2) - \gamma(\mathcal{X}_2))$. In particular, $\mathcal{X}_2$ is neither ordinary or rational by the choice of $\mathcal{X}$ to be a minimal counterexample. From the proof of Proposition 3.1, the case $\mathbf{g}(\mathcal{X}_2) = 1$ cannot occur as $|G_1| > 34(\mathbf{g}(\mathcal{X}_1) + 1)^{3/2}$. Therefore, $\mathbf{g}(\mathcal{X}_2) \geq 2$ and $\mathbf{g}(\mathcal{X}_2) > \gamma(\mathcal{X}_2)$ hold. The factor group $G_2 = G_1/N_2$ is a $\mathbb{K}$ -automorphism group of the quotient curve $\mathcal{X}_2 = \mathcal{X}_1/N_2$, and it has a minimal normal d -subgroup with $d \neq p$, by the choice of N_2 . Define N_3 to be the largest normal, prime to p , subgroup of G_2 . Observe that N_3 must be a proper subgroup of G_2 , otherwise G_2 itself would be a prime to p subgroup of $\text{Aut}(\mathcal{X}_2)$ of order bigger than $34(\mathbf{g}(\mathcal{X}_2) + 1)^{3/2}$ contradicting the classical Hurwitz bound $84(\mathbf{g}(\mathcal{X}_2) - 1)$. Therefore, there exists a (maximal) nontrivial normal p -subgroup N_4 in the factor group $G_3 = G_2/N_3$. Now, the above argument remains valid whenever $G, N_1, G_1, N_2, \mathcal{X}_1, \mathcal{X}_2$ are replaced by $G_2, N_3, G_3, N_4, \mathcal{X}_3, \mathcal{X}_4$ where the quotient curves are $\mathcal{X}_3 = G_2/N_3$ and $\mathcal{X}_4 = G_3/N_4$. In particular, $\mathbf{g}(\mathcal{X}_4) \neq 1$ and $\mathbf{g}(\mathcal{X}_3) - \gamma(\mathcal{X}_3) = |N_4|(\mathbf{g}(\mathcal{X}_4) - \gamma(\mathcal{X}_4))$. Repeating the above argument a finite sharply decreasing sequence $\mathbf{g}(\mathcal{X}_1) > \mathbf{g}(\mathcal{X}_2) > \mathbf{g}(\mathcal{X}_3) > \mathbf{g}(\mathcal{X}_4), \dots$ arises. If this sequence has $n + 1$ members then $\mathbf{g}(\mathcal{X}_n) - \gamma(\mathcal{X}_n) = |N_{n+1}|(\mathbf{g}(\mathcal{X}_{n+1}) - \gamma(\mathcal{X}_{n+1}))$ with $\mathbf{g}(\mathcal{X}_{n+1}) = \gamma(\mathcal{X}_{n+1}) = 0$. Therefore, for some (odd) index $m \leq n$, the curve $\mathcal{X}_m$ for some $m \geq 1$ must be ordinary, a contradiction to choice of $\mathcal{X}$ to be a minimal counterexample.

First we investigate the elliptic case. Since $\mathbf{g}(\mathcal{X}) \geq 2$, the Hurwitz genus formula applied to $\bar{\mathcal{X}}$ ensures that N has a short orbit. Let Γ be a short orbit of G containing a short orbit of N . Since N is a normal subgroup of G , Γ is partitioned into short-orbits $\Sigma_1, \dots, \Sigma_k$ of N each of length $|\Sigma_1|$. Take a point R_i from Σ_i for $i = 1, 2, \dots, k$, and set $\Sigma = \Sigma_1$ and $S = S_1$. With this notation, $|G| = |G_S| |\Gamma| = |G_S| k |\Sigma|$, and the Hurwitz genus formula gives

$$(19) \quad 2\mathbf{g}(\mathcal{X}) - 2 \geq \sum_{i=1}^k |\Sigma_i| (|N_{S_i}| - 1) = k |\Sigma| (|N_S| - 1) \geq +\frac{1}{2} k |\Sigma| |N_S| = \frac{1}{2} |G| \frac{|N_S|}{|G_S|}.$$

Also, the factor group $G_S N / N$ is a subgroup of $\text{Aut}(\bar{\mathcal{X}})$ fixing the the point of $\bar{\mathcal{X}}$ lying under S in the cover $\mathcal{X}|\bar{\mathcal{X}}$. From [3, Theorem 11.94 (ii)],

$$\frac{|G_S N|}{|N|} = \frac{|G_S|}{|G_S \cap N|} = \frac{|G_S|}{|N_S|} \leq 12.$$

This and (19) yield $|G| \leq 48(\mathbf{g}(\mathcal{X}) - 1)$, a contradiction with our hypothesis $34(\mathbf{g}(\mathcal{X}) + 1)^{3/2}$.

Therefore, $\mathcal{X}$ is rational. Thus $\bar{G}$ is isomorphic to a subgroup of $PGL(2, \mathbb{K})$. Since p divides $|G|$ but $|N|$, $\bar{G}$ contains a nontrivial p -subgroup. From the classification of the finite subgroups of $PGL(2, \mathbb{K})$, see [13], either $p = 3$ and $\bar{G} \cong \text{Alt}_4, \text{Sym}_4$, or $\bar{G} = \bar{Q} \rtimes \bar{C}$ where $\bar{Q}$ is a normal p -subgroup and its complement $\bar{C}$ is a cyclic prime to p subgroup and $|\bar{C}|$ divides $|\bar{Q}| - 1$.

If $\bar{G} \cong \text{Alt}_4, \text{Sym}_4$ then $|\bar{G}| \leq 24$ whence $|G| \leq 24|N| \leq 96(\mathbf{g}(\mathcal{X}) + 1)$ as N is abelian. Comparison with our hypothesis $|G| \geq 34(\mathbf{g}(\mathcal{X}) + 1)^{3/2}$ shows that $\mathbf{g}(\mathcal{X}) \leq 6$. For small genera we need a little more. If $|N|$ is prime then $|N| \leq 2\mathbf{g}(\mathcal{X}) + 1$; see [3, Theorem 11.108], and hence $|G| \leq 48(\mathbf{g}(\mathcal{X}) + 1)$ which is inconsistent with $|G| \geq 34(\mathbf{g}(\mathcal{X}) + 1)^{3/2}$. Otherwise, since $p = 3$ and $|N|$ has order a power of prime distinct from p , the bound $|N| \leq 4(\mathbf{g}(\mathcal{X}) + 1)$ with $\mathbf{g}(\mathcal{X}) \leq 6$ is only possible for $(\mathbf{g}(\mathcal{X}), |N|) \in \{(3, 16), (4, 16), (5, 16), (6, 16), (6, 25)\}$. Comparison of $|G| \leq 24|N|$ with $|G| \geq 34(\mathbf{g}(\mathcal{X}) + 1)^{3/2}$ rule out the latter three cases. Furthermore, since N is an elementary abelian group of order 16, $\mathbf{g}(\mathcal{X})$ must be odd by Lemma 2.2. Finally, $\mathbf{g}(\mathcal{X}) = 3, |N| = 16, G/N \cong \text{Sym}_4$ is impossible as Henn's bound $|G| \geq 8\mathbf{g}(\mathcal{X})^3$ implies that $\mathcal{X}$ has zero p -rank, see [3, Theorem 11.127].

Therefore, the case $\bar{G} = \bar{Q} \rtimes \bar{C}$ occurs. Also, $\bar{G}$ fixes a unique place $\bar{P} \in \bar{\mathcal{X}}$. Let Δ be the N -orbits in $\mathcal{X}$ lies over $\bar{P}$ in the cover $\mathcal{X}|\bar{\mathcal{X}}$. We prove that Δ is a long orbit of N . By absurd, the permutation

representation of G on Δ has a nontrivial 1-point stabilizer containing a nontrivial subgroup M of N . Since N is abelian, M is in the kernel. In particular, M is a normal subgroup of G contradicting our choice of N to be minimal.

Take a Sylow p -subgroup Q of G of order $|Q| = p^h$ with $h \geq 1$, and look at the action of Q on Δ . Since $|\Delta| = |N|$ is prime to p , Q fixes a point $P \in \Delta$, that is, $Q = Q_P$. Since $\mathcal{X}$ is an ordinary curve, Q_P and hence Q is elementary abelian; see [8, Theorem 2 (i)] or [3, Theorem 11.74 (iii)]. Therefore, $G_P = Q \rtimes U$ where U is a prime to p cyclic group. Thus

$$(20) \quad |\bar{Q}||\bar{C}||N| = |\bar{G}||N| = |G| = |G_P||\Delta| = |Q||U||\Delta| = |Q||U||N|,$$

whence $|Q| = |\bar{Q}|$ and $|U| = |\bar{C}|$. Consider the subgroup H of G generated by G_P and N . Since Δ is a long N -orbit, $G_P \cap N = \{1\}$. As N is normal in H this implies that $H = N \rtimes G_P = N \rtimes (Q \rtimes U)$ and hence $|H| = |N||Q||U|$ which proves $G = H = N \rtimes (Q \rtimes U)$.

Since $\mathcal{X}$ is rational and $\bar{P}$ is the unique fixed point of nontrivial elements of $\bar{Q}$, each $\bar{Q}$ -orbit other than $\{\bar{P}\}$ is long. Furthermore, $\bar{C}$ fixes a point $\bar{R}$ other than $\bar{P}$ and no nontrivial element of $\bar{C}$ fixes point distinct from $\bar{P}$ and $\bar{R}$. This shows that the $\bar{G}$ -orbit $\bar{\Omega}_1$ of $\bar{R}$ has length $|Q|$. In terms of the action of G on $\mathcal{X}$, there exist as many as $|Q|$ orbits of N , say $\Delta_1, \dots, \Delta_{|Q|}$, whose union Λ is a short G -orbit lying over $\bar{\Omega}_1$ in the cover $\mathcal{X}|\bar{\mathcal{X}}$. Obviously, if at least one of Δ_i is a short N -orbit then so all are.

We show that this actually occur. Since the cover $\mathcal{X}|\bar{\mathcal{X}}$ ramifies, N has some short orbits, and by absurd there exists a short N -orbit Σ is not contained in Λ . Then Σ and Λ are disjoint. Let Γ denote the (short) G -orbit containing Σ . Since N is a normal subgroup of G , Γ is partitioned into N -orbits, say $\Sigma = \Sigma_1, \dots, \Sigma_k$, each of them of the same length $|\Sigma|$. Here $k = |Q||U|$ since the set of points of $\bar{\mathcal{X}}$ lying under these k short N -orbits is a long $\bar{G}$ -orbit. Also, $|N| = |\Sigma_i||N_{R_i}|$ for $i \leq k$ and $R_i \in \Sigma_i$. In particular, $|\Sigma_1| = \Sigma_i$ and $|N_{R_1}| = |B_{R_i}|$. From the Hurwitz genus formula,

$$2g(\mathcal{X}) - 2 \geq -2|N| + \sum_{i=1}^k |\Sigma_i|(|N_{R_i}| - 1) = -2|N| + |Q||U||\Sigma_1|(|N_{R_1}| - 1).$$

Since N_{R_1} is nontrivial, $|N_{R_1}| - 1 \geq \frac{1}{2}|N_{R_1}|$. Therefore,

$$2g(\mathcal{X}) - 2 \geq -2|N| + \frac{1}{2}|Q||U||\Sigma_1||N_{R_1}| = -2|N| + \frac{1}{2}|Q||U||N| = |N|(\frac{1}{2}(|Q||U| - 2)) = \frac{1}{2}|N|(|Q||U| - 4).$$

As $|Q||U| - 4 \geq \frac{1}{2}|Q||U|$ by $|Q||U| \geq 4$, this gives

$$2g(\mathcal{X}) - 2 \geq \frac{1}{4}|N||U||Q| = \frac{1}{4}|G|.$$

But this contradicts our hypothesis $|G| > 34(g(\mathcal{X}) + 1)^{3/2}$.

Therefore, the short orbits of N are exactly $\Delta_1, \dots, \Delta_{|Q|}$. Take a point S_i from Δ_i for $i = 1, \dots, |Q|$. Then N_{S_1} and N_{S_i} are conjugate in G , and hence $|N_{S_1}| = |N_{S_i}|$. From the Hurwitz genus formula applied to N ,

$$2g(\mathcal{X}) - 2 = -2|N| + \sum_{i=1}^{|Q|} |\Delta_i|(|N_{S_i}| - 1) = -2|N| + |Q||\Delta_1|(|N_{S_1}| - 1) \geq -2|N| + \frac{1}{2}|Q||\Delta_1||N_{S_1}|.$$

Since $|N| = |\Delta_1||N_{S_1}|$, this gives $2g(\mathcal{X}) - 2 \geq \frac{1}{2}|N|(|Q| - 4)$ whence $2g(\mathcal{X}) - 2 \geq \frac{1}{4}|N||Q|$ provided that $|Q| \geq 5$. The missing case, $|Q| = 3$, cannot actually occur since in this case $|\bar{C}| = |U| \leq |Q| - 1 = 2$, whence $|G| = |Q||U||N| \leq 6|N| \leq 24(g(\mathcal{X}) + 1)$, a contradiction with $|G| > 34(g(\mathcal{X}) + 1)^{3/2}$. Thus

$$(21) \quad |N||Q| \leq 8(g(\mathcal{X}) - 1).$$

Since $|N||U| < |N||Q|$, this also shows

$$(22) \quad |N||U| < 8(g(\mathcal{X}) - 1).$$

Therefore,

$$|G||N| = |N|^2|U||Q| < 64(g(\mathcal{X}) - 1)^2.$$

Equations (21) and (22) together with our hypothesis $|G| \geq 34(\mathfrak{g}(\mathcal{X}) + 1)^{3/2}$ yield

$$(23) \quad |N| < \frac{64}{34} \sqrt{\mathfrak{g}(\mathcal{X}) - 1}.$$

From (23) and $|G| = |N||Q||U| \geq 34(\mathfrak{g}(\mathcal{X}) + 1)^{3/2}$ we obtain

$$|Q||U| > \frac{34^2}{64}(\mathfrak{g}(\mathcal{X}) - 1) > 18(\mathfrak{g}(\mathcal{X}) - 1)$$

which shows that Lemma 2.1 applies to the subgroup $Q \rtimes U$ of $\text{Aut}(\mathcal{X})$. With the notation in Lemma 2.1, this gives that $Q \rtimes U$ and Q have the same two short orbits, $\Omega_1 = \{P\}$ and Ω_2 . In the cover $\mathcal{X}/\bar{\mathcal{X}}$, the point $\bar{P} \in \bar{\mathcal{X}}$ lying under P is fixed by Q . We prove that Ω_2 is a subset of the N -orbit Δ containing P . For this purpose, it suffices to show that for any point $R \in \Omega_2$, the point $\bar{R} \in \bar{\mathcal{X}}$ lying under R in the cover $\mathcal{X}/\bar{\mathcal{X}}$ coincides with $\bar{P}$. Since Ω_2 is a Q -short orbit, the stabilizer Q_R is nontrivial, and hence Q fixes $\bar{R}$. Since $\mathcal{X}$ is rational, this yields $\bar{P} = \bar{R}$. Therefore, $\Omega_2 \cup \{P\}$ is contained in Δ , and either $\Delta = \Omega_2 \cup \{P\}$ or Δ contains a long Q -orbit. In the latter case, $|U| < |Q| < |N|$, and hence

$$|G|^2 = |N||Q||N||U||Q||U| < |N||Q||N||U||N|^2 \leq \frac{64^2}{34}(\mathfrak{g}(\mathcal{X}) - 1)^3$$

whence $|G| < 34(\mathfrak{g}(\mathcal{X}) + 1)^{3/2}$, a contradiction with our hypothesis. Otherwise $|N| = |\Delta| = 1 + |\Omega_2|$. In particular, $|N|$ is even, and hence it is a power of 2. Also, from the Deuring-Shafarevič Formula, $\mathfrak{g}(\mathcal{X}) - 1 = \gamma(\mathcal{X}) - 1 = -|Q| + 1 + |\Omega_2|$ where $|\Omega_2| \geq 1$ is a power of p . This implies that $\mathfrak{g}(\mathcal{X})$ is also even. Since N is an elementary abelian 2-group, Lemma 2.2 yields that either $|N| = 2$ or $|N| = 4$.

If $|N| = 2$ then Ω_2 consists of a unique point R and $Q \rtimes U$ fixes both points P and R . Since $\Delta = \{P, R\}$, and Δ is a G -orbit, the stabilizer $G_{P,R}$ is an index 2 (normal) subgroup of G . On the other hand, $G_{P,R} = Q \rtimes U$ and hence Q is the unique Sylow p -subgroup of $Q \rtimes U$. Thus Q is a characteristic subgroup of the normal subgroup $G_{P,R}$ of G . But then Q is a normal subgroup of G , a contradiction with our hypothesis.

If $|N| = 4$ then $|\Delta| = 4$ and $p = 3$. The permutation representation of G of degree 4 on Δ contains a 4-cycle induced by N but also a 3-cycle induced by Q . Hence if $K = \ker$ then $G/K \cong \text{Sym}_4$. On the other hand, since both N and Ker are normal subgroups of G , their product NK is normal, as well. Hence NK/K is a normal subgroup of G/K , but this contradicts $G/K \cong \text{Sym}_4$. $\square$

REFERENCES

- [1] M. Giulietti and G. Korchmáros, Garden of curves with many automorphisms, in *Algebraic curves and finite fields*, 93-120, Radon Ser. Comput. Appl. Math., 16, De Gruyter, Berlin, 2014.
- [2] G.B. Gunby, A. Smith and A. Yuan, Irreducible canonical representations in positive characteristic, *Research in Number Theory* (2015) 1:3, 1-25.
- [3] J.W.P. Hirschfeld, G. Korchmáros, and F. Torres, *Algebraic Curves over a Finite Field*, Princeton Series in Applied Mathematics, Princeton (2008).
- [4] H.W. Henn, Funktionenkörper mit großer Automorphismengruppe, *J. Reine Angew. Math.* **302** (1978), 96–115.
- [5] A. Kontogeorgis and V. Rotger, On abelian automorphism groups of Mumford curves, *Bull. London Math. Soc.* **40** (2008), 353-362.
- [6] R. Lidl and H. Niederreiter, *Finite Fields*, Encyclopedia of Mathematics and its Applications **20**, Cambridge Univ. Press, Cambridge, 1997, xiv+755 pp.
- [7] A. Machì, *Groups, An introduction to ideas and methods of the theory of groups*, Unitext, **58**, Springer, Milan, 2012. xiv+371 pp.
- [8] S. Nakajima, p -ranks and automorphism groups of algebraic curves, *Trans. Amer. Math. Soc.* **303** (1987), 595-607.
- [9] J.P. Serre, *Corps locaux*, Hermann, Paris, (1968).
- [10] J.-P. Serre, *Local Fields*, Graduate Texts in Mathematics **67**, Springer, New York, (1979). viii+241 pp.
- [11] H. Stichtenoth, Über die Automorphismengruppe eines algebraischen Funktionenkörpers von Primzahl- charakteristik. II. Ein spezieller Typ von Funktionenkörpern, *Arch. Math.* **24** (1973), 615–631.
- [12] F. Sullivan, p -torsion in the class group of curves with many automorphisms, *Arch. Math.* **26** (1975), 253–261.
- [13] R.C. Valentini and M.L. Madan, A Hauptsatz of L.E. Dickson and Artin-Schreier extensions, *J. Reine Angew. Math.* **318** (1980), 156–177.

Authors' addresses:

Gábor KORCHMÁROS and Maria MONTANUCCI

Dipartimento di Matematica, Informatica ed Economia

Università degli Studi della Basilicata

Contrada Macchia Romana

85100 Potenza (Italy).

E-mail: `gabor.korchmaros@unibas.it` and `maria.montanucci@unibas.it`